

INTRAVENOUS STAT DOSES

18 kg

Affix Patient Label Here

Diluents:

NS = Sodium Chloride 0.9%, G5 = Glucose 5%, WFI =

Water for Injections

● = Central or intraosseus route only

CD = Controlled drug

Maximum doses highlighted in red



Note: Ideal or adjusted body weight should be used where clinically appropriate

Intubation/Induction				
Drug	Dose/kg		Calculated Dose	Dilution
Atracurium besilate	0.5	mg/kg	9 mg	Neat or with NS/G5 to convenient vol.
Atropine sulfate	20	microg/kg	360 microg	Neat or with NS/G5 to convenient vol.
Fentanyl CD	1 - 5	microg/kg	18 - 90 microg	Neat or with NS/G5 to convenient vol.
Ketamine CD	1 - 2	mg/kg	18 - 36 mg	Neat or with NS/G5 to convenient vol.
Midazolam CD	0.1 - 0.2	mg/kg	1.8 - 3.6 mg	Neat or with NS/G5 to convenient vol.
Rocuronium bromide	0.6 - 1	mg/kg	10.8 - 18 mg	Neat or with NS/G5 to convenient vol.
Suxamethonium chloride	2	mg/kg	36 mg	Neat or with NS/G5 to convenient vol.
Thiopental sodium	2 - 5	mg/kg	36 - 90 mg	Reconstitute to 25mg/mL with 20mL NS/G5/WFI

Fluid Bolus	
Dose/kg	Calculated Volume
10mL/kg	180 mL
5mL/kg	90 mL

Maintenance Fluids		
%	Daily Allowance	Rate
100%	1400 mL/day	58.3 mL/hr
75%	1050 mL/day	43.8 mL/hr
50%	700 mL/day	29.2 mL/hr

Cardiac Arrest/Arrhythmias					
Drug	Dose/kg		Calculated Dose	Administer Over	Dilution
Adenosine	0.1 - 0.5	mg/kg	1.8 - 9 mg	2 seconds	Neat or dilute to 0.5mg/mL with 6mL NS
Adrenaline (1:10,000) [100microg/mL]	0.01	mg/kg	0.18 mg	Rapid injection	Dilute to 10mL with NS (max. 1mg/10mL [neat])
Amiodarone hydrochloride cardiac arrest loading dose - arrhythmias	5 - 10	mg/kg	90 - 180 mg	3 minutes 20 minutes - 2 hours	Neat or dilute to 7.5 - 15mg/mL with G5 Dilute to 0.6 - 2.4mg/mL with G5
Calcium <u>GLUCONATE</u> 10% [0.22mmol/mL]	0.11	mmol/kg	1.98 mmol	5 - 10 minutes	Neat or dilute each 1mL with 4mL of NS/GS
Glucose 10%	2	mL/kg	36 mL	2 - 5 minutes	Neat
Magnesium sulfate 50% [500mg/mL]	25 - 50	mg/kg	450 - 900 mg	20 minutes	Dilute each 1mL with 4mL of NS/G5
● Potassium chloride 15% [2mmol/mL] CD	0.4	mmol/kg	7.2 mmol	1 hour	Dilute to 0.5mmol/mL with NS (preferred)/G5
Sodium bicarbonate 8.4% [1mmol/mL]	1	mmol/kg	18 mmol	At least 3 minutes	<2 years: dilute with equal vol. of NS/G5/WFI

Seizures/Intracranial Pressure					
Drug	Dose/kg		Calculated Dose	Administer Over	Dilution
Lorazepam CD	0.1	mg/kg	1.8 mg	Rapid injection over 1 minute	Dilute to 2mg/mL with equal volume of NS/G5
Levetiracetam	40	mg/kg	720 mg	5 minutes	Dilute to 50mg/mL (neonates: 100mg/mL) with NS/G5
Phenobarbital CD	20	mg/kg	360 mg	20 minutes	Dilute to 20mg/mL with WFI
Phenytoin	20	mg/kg	360 mg	20 minutes	Dilute to 5-10mg/mL with NS (use 0.22-0.5micron filter)
Sodium chloride 2.7 - 3%	3 - 5	mL/kg	54 - 90 mL	10 - 15 minutes	Neat (2.7% preferred as pre-made solution)
Mannitol 20%	0.25 - 1	g/kg	22.5 - 90 mL	15 - 30 minutes	Neat (15micron filter recommended)

Asthma					
Drug	Dose/kg		Calculated Dose	Administer Over	Dilution
Magnesium sulfate 50% [500mg/mL]	40	mg/kg	720 mg	Over 20 minutes	Dilute to 20mL with NS
Salbutamol 1-23 months over 2 years	5	microg/kg	90 microg	5-10 minutes	Dilute to 5mL with NS/G5
	15	microg/kg	250 microg	5-10 minutes	Dilute to 5mL with NS/G5

INTRAVENOUS INFUSIONS

18 kg

Affix Patient Label Here

Diluents:

NS = Sodium Chloride 0.9%, G5 = Glucose 5%,

WFI = Water for Injections

● = Central or intraosseus route only

CD = Controlled drug

Maximum doses highlighted in red



Note: Ideal or adjusted body weight should be used where clinically appropriate

Inotropes/Vasoactives						
Infusion	Drug Amount	Diluent	Total Volume	Dose	Rate	
● Adrenaline (1:1000) [1mg/mL]	central	5.4 mg	NS or G5	50 mL	0.1 - 1.5 microg/kg/min	1 mL/hr = 0.1 microg/kg/min
Adrenaline (1:1000) [1mg/mL]	peripheral	5.4 mg	NS or G5	500 mL	0.1 - 1.5 microg/kg/min	1 mL/hr = 0.01 microg/kg/min
● Amiodarone hydrochloride		120 mg	G5	50 mL	5 - 25 microg/kg/min	4.5 mL/hr = 10 microg/kg/min
Dinoprostone (Prostin®)	neonates only					
● Isoprenaline	under 12 years	0.54 mg	G5	50 mL	0.02 - 1 microg/kg/min	1 mL/hr = 0.01 microg/kg/min
	over 12 years	3 mg	G5	50 mL	0.5 - 20 microg/min	1 mL/hr = 1 microg/min
Milrinone	peripheral	10 mg	NS or G5	50 mL	0.25 - 0.75 microg/kg/min	2.7 mL/hr = 0.5 microg/kg/min
● Noradrenaline	central	5.4 mg	NS or G5	50 mL	0.1 - 1.5 microg/kg/min	1 mL/hr = 0.1 microg/kg/min
Noradrenaline	peripheral	5.4 mg	NS or G5	500 mL	0.1 - 1.5 microg/kg/min	1 mL/hr = 0.01 microg/kg/min
● Vasopressin (argipressin)		18 units	NS or G5	50 mL	0.0003 - 0.002 units/kg/min	1 mL/hr = 0.0003 units/kg/min
Sedatives/Analgesics/Muscle Relaxants						
Infusion	Drug Amount	Diluent	Total Volume	Dose	Rate	
Fentanyl CD	900 microg	NS or G5	50 mL	1 - 10 microg/kg/hr	1 mL/hr =	1 microg/kg/hr
● Ketamine CD	central	540 mg	NS or G5	50 mL	10 - 45 microg/kg/min	1 mL/hr = 10 microg/kg/min
Ketamine CD	peripheral	500 mg	Neat	50 mL	10 - 45 microg/kg/min	1.62 mL/hr = 15 microg/kg/min
Morphine CD	18 mg	NS or G5	50 mL	5 - 60 microg/kg/hr	1 mL/hr =	20 microg/kg/hr
Midazolam CD	54 mg	NS or G5	50 mL	30 - 300 microg/kg/hr	1 mL/hr =	60 microg/kg/hr
Propofol 1%	maximum 12 hours	500 mg	Neat	50 mL	1 - 4 mg/kg/hr	1.8 mL/hr = 1 mg/kg/hr
Rocuronium bromide	500 mg	Neat	50 mL	300 - 1000 microg/kg/hr	1.8 mL/hr =	1000 microg/kg/hr
Bronchodilators						
Infusion	Drug Amount	Diluent	Total Volume	Dose	Rate	
Aminophylline	loading dose	500 mg	NS	500 mL	108 mg (6mg/kg)	108 mL over 20 minutes
	followed by	500 mg	NS	500 mL	1 mg/kg/hr	18 mL/hr = 1 mg/kg/hr
	under 12 years	500 mg	NS	500 mL	0.5 - 0.7 mg/kg/hr	9 mL/hr = 0.5 mg/kg/hr
	over 12 years					
Magnesium sulphate 50% [500mg/mL]	5 g	G5	50 mL	50 mg/kg/hr	9 mL/hr	over 4 hours
Salbutamol	peripheral	10 mg	NS or G5	50 mL	0.5-1 microg/kg/min	2.7 mL/hr = 0.5 microg/kg/min
	monitor for extravasation				maximum 20microg/min	

Disclaimer: All medicines to be given INTRAVENOUSLY unless otherwise stated. It is the responsibility of the clinician to ensure drugs are used appropriately according to the clinical situation and doses double checked. NWTS/ODN does not accept any liability. Use of these monographs is at the clinician's own risk.