

INTRAVENOUS STAT DOSES

22 kg

Diluents:

NS = Sodium Chloride 0.9%, G5 = Glucose 5%, WFI = Water for Injections

• = Central or intraosseus route only

CD = Controlled drug

Maximum doses highlighted in red



Note: Ideal or adjusted body weight should be used where clinically appropriate

Intubation/Induction

Drug	Dose/kg	Calculated Dose	Dilution
Atracurium besilate	0.5 mg/kg	11 mg	Neat or with NS/G5 to convenient vol.
Atropine sulfate	20 microg/kg	440 microg	Neat or with NS/G5 to convenient vol.
Fentanyl CD	1 - 5 microg/kg	22 - 110 microg	Neat or with NS/G5 to convenient vol.
Ketamine CD	1 - 2 mg/kg	22 - 44 mg	Neat or with NS/G5 to convenient vol.
Midazolam CD	0.1 - 0.2 mg/kg	2.2 - 4.4 mg	Neat or with NS/G5 to convenient vol.
Rocuronium bromide	0.6 - 1 mg/kg	13.2 - 22 mg	Neat or with NS/G5 to convenient vol.
Suxamethonium chloride	2 mg/kg	44 mg	Neat or with NS/G5 to convenient vol.
Thiopental sodium	2 - 5 mg/kg	44 - 110 mg	Reconstitute to 25mg/mL with 20mL NS/G5/WFI

Fluid Bolus

Dose/kg	Calculated Volume
10mL/kg	220 mL
5mL/kg	110 mL

Maintenance Fluids

%	Daily Allowance	Rate
100%	1540 mL/day	64.2 mL/hr
75%	1155 mL/day	48.1 mL/hr
50%	770 mL/day	32.1 mL/hr

Cardiac Arrest/Arrhythmias

Drug	Dose/kg	Calculated Dose	Administer Over	Dilution
Adenosine	0.1 - 0.5 mg/kg	2.2 - 11 mg	2 seconds	Neat or dilute to 0.5mg/mL with 6mL NS
Adrenaline (1:10,000) [100microg/mL]	0.01 mg/kg	0.22 mg	Rapid injection	Neat
Amiodarone hydrochloride cardiac arrest loading dose - arrhythmias	5 mg/kg 5 - 10 mg/kg	110 mg 110 - 220 mg	3 minutes 20 minutes - 2 hours	Neat or dilute to 7.5 - 15mg/mL with G5 Dilute to 0.6 - 2.4mg/mL with G5
Calcium GLUCONATE 10% [0.22mmol/mL]	0.11 mmol/kg	2.42 mmol	5 - 10 minutes	Neat or dilute each 1mL with 4mL of NS/G5
Glucose 10%	2 mL/kg	44 mL	2 - 5 minutes	Neat
Magnesium sulfate 50% [500mg/mL]	25 - 50 mg/kg	550 - 1100 mg	20 minutes	Dilute each 1mL with 4mL of NS/G5
• Potassium chloride 15% [2mmol/mL] CD	0.4 mmol/kg	8.8 mmol	1 hour	Dilute to 0.5mmol/mL with NS (preferred)/G5
Sodium bicarbonate 8.4% [1mmol/mL]	1 mmol/kg	22 mmol	At least 3 minutes	<2 years: dilute with equal vol. of NS/G5/WFI

Seizures/Intracranial Pressure

Drug	Dose/kg	Calculated Dose	Administer Over	Dilution
Lorazepam CD	0.1 mg/kg	2.2 mg	Rapid injection over 1 minute	Dilute to 2mg/mL with equal volume of NS/G5
Levetiracetam	40 mg/kg	880 mg	5 minutes	Dilute to 50mg/mL (neonates: 100mg/mL) with NS/G5
Phenobarbital CD	20 mg/kg	440 mg	20 minutes	Dilute to 20mg/mL with WFI
Phenytoin	20 mg/kg	440 mg	20 minutes	Dilute to 5-10mg/mL with NS (use 0.22-0.5micron filter)
Sodium chloride 2.7 - 3%	3 - 5 mL/kg	66 - 110 mL	10 - 15 minutes	Neat (2.7% preferred as pre-made solution)
Mannitol 20%	0.25 - 1 g/kg	27.5 - 110 mL	15 - 30 minutes	Neat (15micron filter recommended)

Asthma

Drug	Dose/kg	Calculated Dose	Administer Over	Dilution
Magnesium sulfate 50% [500mg/mL]	40 mg/kg	880 mg	Over 20 minutes	Dilute to 20mL with NS
Salbutamol 1-23 months over 2 years	5 microg/kg 15 microg/kg	110 microg 250 microg	5-10 minutes 5-10 minutes	Dilute to 5mL with NS/G5 Dilute to 5mL with NS/G5

INTRAVENOUS INFUSIONS

22 kg

Affix Patient Label Here

Diluents:

NS = Sodium Chloride 0.9%, G5 = Glucose 5%,
WFI = Water for Injections

● = Central or intraosseus route only

CD = Controlled drug

Maximum doses highlighted in red



Note: Ideal or adjusted body weight should be used where clinically appropriate

Inotropes/Vasoactives

Infusion	Drug Amount	Diluent	Total Volume	Dose	Rate
● Adrenaline (1:1000) [1mg/mL] central	6.6 mg	NS or G5	50 mL	0.1 - 1.5 microg/kg/min	1 mL/hr = 0.1 microg/kg/min
Adrenaline (1:1000) [1mg/mL] peripheral	6.6 mg	NS or G5	500 mL	0.1 - 1.5 microg/kg/min	10 mL/hr = 0.1 microg/kg/min
● Amiodarone hydrochloride	120 mg	G5	50 mL	5 - 25 microg/kg/min	5.5 mL/hr = 10 microg/kg/min
Dinoprostone (Prostin®) neonates only					
● Isoprenaline under 12 years	0.66 mg	G5	50 mL	0.02 - 1 microg/kg/min	10 mL/hr = 0.1 microg/kg/min
over 12 years	3 mg	G5	50 mL	0.5 - 20 microg/min	1 mL/hr = 1 microg/min
Milrinone peripheral	10 mg	NS or G5	50 mL	0.25 - 0.75 microg/kg/min	3.3 mL/hr = 0.5 microg/kg/min
● Noradrenaline central	6.6 mg	NS or G5	50 mL	0.1 - 1.5 microg/kg/min	1 mL/hr = 0.1 microg/kg/min
Noradrenaline peripheral	6.6 mg	NS or G5	500 mL	0.1 - 1.5 microg/kg/min	10 mL/hr = 0.1 microg/kg/min
● Vasopressin (argipressin)	22 units	NS or G5	50 mL	0.0003 - 0.002 units/kg/min	1 mL/hr = 0.0003 units/kg/min

Sedatives/Analgesics/Muscle Relaxants

Infusion	Drug Amount	Diluent	Total Volume	Dose	Rate
Fentanyl CD	1100 microg	NS or G5	50 mL	1 - 10 microg/kg/hr	1 mL/hr = 1 microg/kg/hr
● Ketamine CD central	660 mg	NS or G5	50 mL	10 - 45 microg/kg/min	1 mL/hr = 10 microg/kg/min
Ketamine CD peripheral	500 mg	Neat	50 mL	10 - 45 microg/kg/min	1.98 mL/hr = 15 microg/kg/min
Morphine CD	22 mg	NS or G5	50 mL	5 - 60 microg/kg/hr	1 mL/hr = 20 microg/kg/hr
Midazolam CD	66 mg	NS or G5	50 mL	30 - 300 microg/kg/hr	1 mL/hr = 60 microg/kg/hr
Propofol 1% maximum 12 hours	500 mg	Neat	50 mL	1 - 4 mg/kg/hr	2.2 mL/hr = 1 mg/kg/hr
Rocuronium bromide	500 mg	Neat	50 mL	300 - 1000 microg/kg/hr	2.2 mL/hr = 1000 microg/kg/hr

Bronchodilators

Infusion	Drug Amount	Diluent	Total Volume	Dose	Rate
Aminophylline loading dose	500 mg	NS	500 mL	132 mg (6mg/kg)	132 mL over 20 minutes
followed by under 12 years	500 mg	NS	500 mL	1 mg/kg/hr	22 mL/hr = 1 mg/kg/hr
over 12 years	500 mg	NS	500 mL	0.5 - 0.7 mg/kg/hr	11 mL/hr = 0.5 mg/kg/hr
Magnesium sulphate 50% [500mg/mL]	5 g	G5	50 mL	50 mg/kg/hr	11 mL/hr over 4 hours
Salbutamol peripheral	10 mg	NS or G5	50 mL	0.5-1 microg/kg/min	3.3 mL/hr = 0.5 microg/kg/min
monitor for extravasation				maximum 20microg/min	

Disclaimer: All medicines to be given INTRAVENOUSLY unless otherwise stated. It is the responsibility of the clinician to ensure drugs are used appropriately according to the clinical situation and doses double checked. NWTS/ODN does not accept any liability. Use of these monographs is at the clinician's own risk. Due to stock shortages and regional variation in available preparations, not every dose can be rounded to a measurable volume - clinical judgement is required to determine the most appropriate dose.

Version 1.2 (updated December 2025), Ratified by: MFT Paediatric Medicines Management Committee (January 2025) Review date: January 2028