

INTRAVENOUS STAT DOSES

55 kg

Diluents:

NS = Sodium Chloride 0.9%, G5 = Glucose 5%, WFI = Water for Injections

• = Central or intraosseus route only

CD = Controlled drug

Maximum doses highlighted in red



Note: Ideal or adjusted body weight should be used where clinically appropriate

Intubation/Induction				
Drug	Dose/kg		Calculated Dose	Dilution
Atracurium besilate	0.5	mg/kg	27.5 mg	Neat or with NS/G5 to convenient vol.
Atropine sulfate	20	microg/kg	600 microg	Neat or with NS/G5 to convenient vol.
Fentanyl CD	1 - 5	microg/kg	55 - 200 microg	Neat or with NS/G5 to convenient vol.
Ketamine CD	1 - 2	mg/kg	55 - 110 mg	Neat or with NS/G5 to convenient vol.
Midazolam CD	0.1 - 0.2	mg/kg	5 mg	Neat or with NS/G5 to convenient vol.
Rocuronium bromide	0.6 - 1	mg/kg	33 - 55 mg	Neat or with NS/G5 to convenient vol.
Suxamethonium chloride	2	mg/kg	100 mg	Neat or with NS/G5 to convenient vol.
Thiopental sodium	2 - 5	mg/kg	110 - 275 mg	Reconstitute to 25mg/mL with 20mL NS/G5/WFI

Fluid Bolus	
Dose/kg	Calculated Volume
10mL/kg	550 mL
5mL/kg	275 mL

Maintenance Fluids		
%	Daily Allowance	Rate
100%	2000 mL/day	83.3 mL/hr
75%	1500 mL/day	62.5 mL/hr
50%	1000 mL/day	41.7 mL/hr

Cardiac Arrest/Arrhythmias					
Drug	Dose/kg		Calculated Dose	Administer Over	Dilution
Adenosine	0.1 - 0.5	mg/kg	3 - 12 mg	2 seconds	Neat or dilute to 0.5mg/mL with 6mL NS
Adrenaline (1:10,000) [100microg/mL]	0.01	mg/kg	0.55 mg	Rapid injection	Neat
Amiodarone hydrochloride cardiac arrest loading dose - arrhythmias	5 - 10	mg/kg	275 - 300 mg	3 minutes 20 minutes - 2 hours	Neat or dilute to 7.5 - 15mg/mL with G5 Dilute to 0.6 - 2.4mg/mL with G5
Calcium GLUCONATE 10% [0.22mmol/mL]	0.11	mmol/kg	6.05 mmol	5 - 10 minutes	Neat or dilute each 1mL with 4mL of NS/G5
Glucose 10%	2	mL/kg	110 mL	2 - 5 minutes	Neat
Magnesium sulfate 50% [500mg/mL]	25 - 50	mg/kg	1375 - 2000 mg	20 minutes	Dilute each 1mL with 4mL of NS/G5
• Potassium chloride 15% [2mmol/mL] CD	0.4	mmol/kg	20 mmol	1 hour	Dilute to 0.5mmol/mL with NS (preferred)/G5
Sodium bicarbonate 8.4% [1mmol/mL]	1	mmol/kg	50 mmol	Rate of 10mmol/min	<2 years: dilute with equal vol. of NS/G5/WFI

Seizures/Intracranial Pressure					
Drug	Dose/kg		Calculated Dose	Administer Over	Dilution
Lorazepam CD	0.1	mg/kg	4 mg	Rapid injection over 1 minute	Dilute to 2mg/mL with equal volume of NS/G5
Levetiracetam	40	mg/kg	2.2 g	5 minutes	Dilute to 50mg/mL (neonates: 100mg/mL) with NS/G5
Phenobarbital CD	20	mg/kg	1 g	20 minutes	Dilute to 20mg/mL with WFI
Phenytoin	20	mg/kg	1.1 g	Rate of 50mg/min	Dilute to 5-10mg/mL with NS (use 0.22-0.5micron filter)
Sodium chloride 2.7 - 3%	3 - 5	mL/kg	165 - 275 mL	10 - 15 minutes	Neat (2.7% preferred as pre-made solution)
Mannitol 20%	0.25 - 1	g/kg	68.8 - 275 mL	15 - 30 minutes	Neat (15micron filter recommended)

Asthma					
Drug	Dose/kg		Calculated Dose	Administer Over	Dilution
Magnesium sulfate 50% [500mg/mL]	40	mg/kg	2 g	Over 20 minutes	Dilute to 20mL with NS
Salbutamol 1-23 months over 2 years	5	microg/kg	250 microg	5-10 minutes	Dilute to 5mL with NS/G5
	15	microg/kg	250 microg	5-10 minutes	Dilute to 5mL with NS/G5

INTRAVENOUS INFUSIONS

55 kg

Affix Patient Label Here

Diluents:

NS = Sodium Chloride 0.9%, G5 = Glucose 5%,
WFI = Water for Injections

● = Central or intraosseous route only

CD = Controlled drug

Maximum doses highlighted in red



Note: Ideal or adjusted body weight should be used where clinically appropriate

Inotropes/Vasoactives						
Infusion		Drug Amount	Diluent	Total Volume	Dose	Rate
● Adrenaline (1:1000) [1mg/mL]	central	16 mg	NS or G5	50 mL	0.1 - 1.5 microg/kg/min	1.03 mL/hr = 0.1 microg/kg/min
Adrenaline (1:1000) [1mg/mL]	peripheral	16 mg	NS or G5	500 mL	0.1 - 1.5 microg/kg/min	10.31 mL/hr = 0.1 microg/kg/min
● Amiodarone hydrochloride		120 mg	G5	50 mL	5 - 25 microg/kg/min	13.75 mL/hr = 10 microg/kg/min
Dinoprostone (Prostin®)	neonates only					
● Isoprenaline	under 12 years	1.65 mg	G5	50 mL	0.02 - 1 microg/kg/min	10 mL/hr = 0.1 microg/kg/min
	over 12 years	3 mg	G5	50 mL	0.5 - 20 microg/min	1 mL/hr = 1 microg/min
Milrinone	peripheral	10 mg	NS or G5	50 mL	0.25 - 0.75 microg/kg/min	8.25 mL/hr = 0.5 microg/kg/min
● Noradrenaline	central	16 mg	NS or G5	50 mL	0.1 - 1.5 microg/kg/min	1.03 mL/hr = 0.1 microg/kg/min
Noradrenaline	peripheral	8 mg	NS or G5	500 mL	0.1 - 1.5 microg/kg/min	20.63 mL/hr = 0.1 microg/kg/min
● Vasopressin (argipressin)		50 units	NS or G5	50 mL	0.0003 - 0.002 units/kg/min	0.99 mL/hr = 0.0003 units/kg/min
Sedatives/Analgesics/Muscle Relaxants						
Infusion		Drug Amount	Diluent	Total Volume	Dose	Rate
Fentanyl CD		2500 microg	Neat	50 mL	1 - 10 microg/kg/hr	1.1 mL/hr = 1 microg/kg/hr
● Ketamine CD	central	1650 mg	NS or G5	50 mL	10 - 45 microg/kg/min	1 mL/hr = 10 microg/kg/min
Ketamine CD	peripheral	500 mg	Neat	50 mL	10 - 45 microg/kg/min	4.95 mL/hr = 15 microg/kg/min
Morphine CD		50 mg	NS or G5	50 mL	5 - 60 microg/kg/hr	1.1 mL/hr = 20 microg/kg/hr
Midazolam CD		100 mg	NS or G5 or Neat	50 mL	0 - 10 mg/hr	1 mL/hr = 2 mg/hr
Propofol 1%	maximum 12 hours	500 mg	Neat	50 mL	55 - 200 mg/hr	10 mL/hr = 100 mg/hr
Rocuronium bromide		500 mg	Neat	50 mL	300 - 1000 microg/kg/hr	5.5 mL/hr = 1000 microg/kg/hr
Bronchodilators						
Infusion		Drug Amount	Diluent	Total Volume	Dose	Rate
Aminophylline	loading dose	500 mg	NS	500 mL	330 mg (6mg/kg)	330 mL over 20 minutes
	followed by					
	under 12 years	500 mg	NS	500 mL	1 mg/kg/hr	55 mL/hr = 1 mg/kg/hr
	over 12 years	500 mg	NS	500 mL	0.5 - 0.7 mg/kg/hr	27.5 mL/hr = 0.5 mg/kg/hr
Magnesium sulphate 50% [500mg/mL]		5 g	G5	50 mL	2 g/hr	20 mL/hr over 4 hours
Salbutamol	peripheral	10 mg	NS or G5	50 mL	20 microg/min	6 mL/hr = 20 microg/min
monitor for extravasation						

Disclaimer: All medicines to be given INTRAVENOUSLY unless otherwise stated. It is the responsibility of the clinician to ensure drugs are used appropriately according to the clinical situation and doses double checked. NWTS/ODN does not accept any liability. Use of these monographs is at the clinician's own risk. Due to stock shortages and regional variation in available preparations, not every dose can be rounded to a measurable volume - clinical judgement is required to determine the most appropriate dose.

Version 1.2 (updated December 2025), Ratified by: MFT Paediatric Medicines Management Committee (January 2025) Review date: January 2028